

Naloxone Medical History

Name _____ CRA # _____

Address _____

Year of Birth _____ Age _____ Race _____ Male Female

Age first use of opiates _____ Age first needle use _____

Which opiates using? Heroin methadone other _____

How much? _____ Daily _____ Occasional _____

In the past year, how many times have you gone 3 or more days without any opiates and why?

(rehab, jail, etc) _____ Last time? _____

In the past six months, which other drugs have you used regularly (> 1-2 times/month)?

Cocaine Y N Valium/Xanax/Ativan/Klonopin Y N

Alcohol Y N Speed (including crystal) Y N

Clonidine Y N PCP Y N

times you have OD'd? _____ last time? _____ On what? _____

times witness OD? _____ # taken to hospital? _____ # died? _____

times you had an abscess? _____ last time? _____

Are you taking any medications? Y N If so, what? _____

Date of last HIV test _____ Date of last HCV test _____ Want to be tested? _____

How many times have you been to an emergency room (for any reason) in the last

two years? _____ How many times have you been admitted (stayed overnight) at a

hospital in the last two years, and for what? _____

participant
Initials

Naloxone Checklist & Order to Dispense

OD prevention techniques

Educate those you use with! Purity testing, tie-release, gradual injecting

Risk factors: mixing drugs, abstinence, use of naltrexone (Trexan™ or ReVia™)

Signs of Overdose

Sleepiness; Slower breathing; Non-responsive to verbal/touch: Turning blue

Call for help → 911 / 311 / Private ambulance / Yell!

A & B of Life -- no one dies from opiate overdose if they can be kept breathing!!!

Airway Remove gum, food, anything from mouth; put in Recovery Position if Breathing

If they stop or have slowed breathing → YOU MUST BREATHE FOR THEM

Rescue Breathing

On back, lift up on back of neck -- tip chin to open airway

Remember to clear mouth and pinch off nose, seal your mouth over theirs

Two quick breaths to begin, then 1 breath every five seconds

Naloxone - store away from light and at room temperature

keep naloxone and IM (1 - 1 1/2") needles with your works

1-2 cc or ml into shoulder; butt cheek; thigh – ***Use long IM needle (1 - 1 1/2 inch needle)***

Breathe for them until it works

If not working in 5 minutes, try another dose

If 2nd dose doesn't work in 5 minutes, something else is wrong → GET HELP

Return of overdose – naloxone lasts 30-90 minutes in the blood

Heroin overdose may last 2 hours

Methadone overdose may last 24 hours!!! → get to hospital !!

Multiple OD (alcohol, benzos, cocaine) → get to hospital !!

SCARE ME pocket card and Partner Checklist

Distribute Naloxone 0.4 mg/ml, 10ml vial Lot # _____ Exp date _____
Distribute at least four 3-10cc syringes with 18-23g 1-1.5 inch needles with each vial of naloxone.

Outreach Worker

Date

Prescriber

Date